

DATA SHEET
CENTEON MAX®
(WP Formulation)
Microbial plant biostimulant.
Polvo mojable (WP)

CENTEON MAX® WP is a biological biostimulant composed of *Trichoderma virens* and *Bacillus subtilis*, formulated as a wettable powder with wide applicability on seeds and plants, improving vegetative development. Activates root development. Its application helps to overcome deficiencies in plant growth.

Approved by S.A.G. for Use in National Organic Agriculture

Formulation:

COMPOSITION	(%w/w)
<i>Trichoderma virens</i> cepa	7*10 ⁷ CFU/g
Luito <i>Bacillus subtilis</i> cepa N5	3*10 ⁷ CFU/g
Coformulantes c.s.p.	94% w/w (940 g/kg)

INSTRUCTIONS FOR USE

CROP	DOSE	TIME OF APPLICATION
Berries	1 to 2 kg/ha (200 g/100L of water for drainage use)	Apply in plantations, to the ground or by immersion of plants or with irrigation. In established plants, it allows applications in budding, development, harvesting, and post-harvesting. through irrigation or sprinkling to the soil. Consider per planting hole 3 - 5 g/plant as Dusting.
Fruit Trees & Vines	1.5 to 2kg/ha (500 g/100L of water for Use in Drenching)	Apply in plantations, to the ground or by immersion of plants or with irrigation. In established plants, it allows applications in budding, development, harvesting, and post-harvesting, through irrigation or sprinkling to the soil. Consider per planting hole 3 - 5 g/plant as Dusting.
Crops (Cereals; Oilseeds; Pope)	1 to 2 kg/ha	Apply from sowing to harvest to the soil by spraying.
Vegetables	2 to 5 g/L water	In storage, applications from sowing to transplanting.
	1 to 2 kg/ha	Application in the field, pre-transplant to the substrate or by immersion of seedlings. Post-transplantation via irrigation or drainage, at all stages of the crop.
Forest	2 to 5 g/L water	Nursery applications from sowing to transplanting.
	1 to 2 kg/ha	Application in the field, pre-transplant to the substrate or by immersion of seedlings. Post-transplants road irrigation or draining, at all stages of cultivation.
Flowers & Ornamentals	2 to 5 g/L of water	Nursery application from sowing to transplanting.
	1 to 2 kg/ha	Application in the field, pre-transplant to substrate or by immersion of seedlings. Post-transplants road irrigation or draining, at all stages of cultivation.

Number of applications: There is no limit on the number of applications per season

Preparation of the mixture: Shake in a bucket with little water the necessary amount of **CENTEON MAX® WP** to be used. Fill the pump tank halfway with water. Empty the contents of the bucket into the pond and fill the volume of water by stirring.

How to apply: Apply via irrigation or drainage. Use a wetting volume that allows the correct coverage of plants, soil and/or substrate.

Compatibility: Incompatibility for joint applications with cupric compounds, fungicides and antibiotics. Do not mix **CENTEON MAX® WP** with pesticides and agrochemicals without first performing a compatibility test or on the manufacturer's instructions. Check with the manufacturer's compatibility list.

Phytotoxicity: There is no phytotoxicity in recommended crops.

Storage: It should be stored in a cool, dry environment for a maximum of 15 months. Keep in its original container, with its respective label.

Waiting Period: There are no waiting days or re-entry restrictions.

Use in organic farming:

**Product suitable for use in Organic Farming in accordance with:
Regulations (EU) No. 2018/848 and 2021/1165, NOP Regulation and
the JAS Regulation.
Control ECOCERT F-32600**



**It is listed on OMRI®.
Approved by S.A.G. for Use in National Organic Agriculture**

EMERGENCY TELEPHONES:

CITUC TOXICOLOGICAL EMERGENCIES PHONE: +56-2-26353800

CITUC CHEMICAL EMERGENCIES PHONE: +56-2-22473600 - 24-HOUR

SERVICE BIO SUPPLIES NATIVA SPA (+56-71) 2970696/2970698.



*Bio Insumos Nativa SpA
Dirección Parcela Antihue Lote 4B 2, Maule-Talca, Chile
Dirección Postal Casilla 6 Maule-Talca, Chile
Fono +56 71 2970696
www.bionativa.cl*